

## Clinical Guideline for: Antacid Prophylaxis for Obstetric Anaesthesia

<b>For Use in:</b>	All Maternity areas
<b>By:</b>	Medical and Midwifery Staff
<b>For:</b>	Patients
<b>Division responsible for document:</b>	Women and Children's Services
<b>Key words:</b>	Obstetric anaesthesia, aspiration, antacid
<b>Name of document author:</b>	Jonathon Francis
<b>Job title of document author:</b>	Consultant
<b>Name of document author's Line Manager:</b>	Anna Haestier
<b>Job title of author's Line Manager:</b>	Chief of Maternity Service
<b>Supported by:</b>	Charles Bircher
<b>Assessed and approved by the:</b>	Maternity Guidelines Committee <b>If approved by committee or Governance Lead Chair's Action; tick here</b> ✓
<b>Date of approval:</b>	24 <sup>th</sup> September 2021
<b>Ratified by or reported as approved to (if applicable):</b>	Clinical Standards Group and Clinical Safety and Effectiveness sub-board
<b>To be reviewed before:</b> This document remains current after this date but will be under review	24 <sup>th</sup> September 2024
<b>To be reviewed by:</b>	Jon Francis
<b>Reference and / or Trust Docs ID No:</b>	825
<b>Version No:</b>	3.2
<b>Compliance links:</b> <i>(is there any NICE related to guidance)</i>	NICE Clinical Guideline: Intrapartum Care – care of healthy women and their babies during childbirth. RCOG Press; September 2007
<b>If Yes - does the strategy/policy deviate from the recommendations of NICE? If so why?</b>	No

# Clinical Guideline for: Antacid Prophylaxis for Obstetric Anaesthesia

## Version and Document Control:

Version Number	Date of Update	Change Description	Author
3.2	24/09/2021	Replacement of Ranitidine with Omeprazole	Jon Francis

### This is a Controlled Document

Printed copies of this document may not be up to date. Please check the hospital intranet for the latest version and destroy all previous versions.

# Clinical Guideline for: Antacid Prophylaxis for Obstetric Anaesthesia

## Background

Pulmonary aspiration of gastric contents is a significant cause of death accompanying anaesthesia in obstetric practice. The incidence of Mendelson's syndrome is reduced by antacid prophylaxis for women having an emergency or elective operative delivery.

To reduce the risk of aspiration pneumonitis, national guidelines recommend that women having a CS (or at high risk of having a CS) should be offered antacid drugs to reduce gastric volumes and acidity.

Patients having an elective caesarean section under regional anaesthesia or general anaesthesia. These patients should be prescribed 40mg Omeprazole at 10pm the night before their CS and a further 40mg Omeprazole at 6am on the morning of surgery.

Patients in labour with a high risk of needing a general anaesthetic.

Patients meeting the following criteria should be given 40mg Omeprazole 12 hourly by mouth:

- Suspected fetal compromise – e.g. Abnormal CTG or FBS, fresh meconium
- Known/suspected placental insufficiency
- Slow progress in labour
- Previous uterine scar
- Multiple pregnancy
- Breech presentation
- People with diabetes
- Pregnancy induced hypertension/PET
- Induction of labour with oxytocin
- Known anaesthetic problem

In addition M/3 sodium citrate 30mL should be given in the anaesthetic room to any woman who is going to have a general anaesthetic.

If a woman in labour has not had oral omeprazole please consult with the on-call anaesthetist

# Clinical Guideline for: Antacid Prophylaxis for Obstetric Anaesthesia

## References

1. Gillett GB, Watson JD, Langford RM. Ranitidine and single-dose antacid therapy as a prophylaxis against acid aspiration syndrome in obstetric practice. *Anaesthesia* 1984;39:638-644
2. National Collaborating Centre for Women's and Children's Health. Caesarean section. Clinical Guideline. London: RCOG Press; 2004.
3. Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer – 2003-2005. The 7th report of the Confidential Enquiries into Maternal Deaths in the UK. RCOG Press, London, December 2007.
4. Sweeney B. and Wright I. The use of antacids as a prophylaxis against Mendelson's syndrome in the United Kingdom. A survey. *Anaesthesia* 1986;41:419-422
5. NICE Clinical Guideline: Intrapartum Care – care of healthy women and their babies during childbirth. RCOG Press; September 2007