

## Clinical Guideline: Umbilical Venous Catheter Insertion

**Authors:** Katie Cullum (ANNP)  
Dr Paul Clarke (Consultant Neonatologist)

**For use in:** EoE Neonatal Units  
Guidance specific to the care of neonatal patients.

**Used by:**

**Key Words:** UVC, umbilical venous catheter insertion, indications, complications, position

**Review due:** December 2023

**Registration No:** NEO-ODN-2020-9

**Approved by:**

Neonatal Clinical Oversight Group	
Clinical Lead Matthew James	<b>Matthew James</b>

**Ratified by Clinical Oversight Group:**

<b>Date of meeting</b>	<b>10<sup>th</sup> December 2020</b>
------------------------	--------------------------------------

**Audit Standards:**

**Audit points**

## 1) Background

The use of umbilical venous catheters (UVCs) is an essential part of neonatal care allowing delivery of intravenous fluids, nutrition, and medication. However, the use of UVCs is associated with a number of potential complications. Whilst catheter-associated infection is well recognised, extravasation into a body cavity is less common but potentially fatal if tamponade ensues. The British Association for Perinatal Medicine has recently published a framework for practice to reduce harm and improve safety in babies needing central venous catheters.<sup>1</sup> The present guideline for UVC insertion incorporates key practice points highlighted by the BAPM expert working group to reduce the risk and complications arising from this procedure.

## 2) Objective

To ensure the safe insertion of an umbilical venous catheter.

## 3) Indications<sup>2</sup>

- Central venous access for low birth weight infants to avoid multiple peripheral cannulation
- Delivery of drugs and parenteral nutrition
- Emergency vascular access for resuscitation of infants at birth
- Exchange transfusion

## 4) Contraindications<sup>2</sup>

- Abnormalities of the abdominal wall
- Necrotising enterocolitis
- Peritonitis

## 5) Complications<sup>1</sup>

- Sepsis<sup>4,5</sup>
- Embolism<sup>3</sup>
- Venous thrombosis<sup>6</sup>
- Pericardial effusion<sup>7,8</sup>
- Pleural effusion<sup>9</sup>
- Portal hypertension
- Displacement leading to blood loss
- Breakage of catheter on removal<sup>10</sup>
- Intra-abdominal extravasation<sup>1</sup>

## 6) Mispositioned catheters

- Catheters placed in the heart can cause pericardial effusion, cardiac tamponade, endocarditis, arrhythmias, and death<sup>1,8,11,12,13,14,15</sup>
- Catheters placed in the portal system are associated with necrotising enterocolitis, perforation of colon and hepatic necrosis<sup>15,16</sup>.
- Catheters in a low position (pre-hepatic) may be at increased risk of hepatic necrosis and intra-abdominal extravasation<sup>1</sup>

## 7) Key notes (based upon the recommendations of the 2018 BAPM Framework<sup>1</sup>)

- A umbilical venous catheter (UVC) tip should lie outside the cardiac silhouette ideally at T8-T9.
- A UVC with tip sited at or below T10 should only be used in the short term (if considered essential) as this site carries a significantly higher risk of extravasation
- Staff inserting UVCs have a responsibility to ensure they maintain their competence and should be familiar with the equipment and procedures used for catheter insertion.
- In each situation where a UVC is required an assessment should be made as to who is the most appropriate person to undertake the procedure.
- A Central Catheter Care bundle should be used to covers all aspects of insertion, use and on-going management of the UVC.
- Staff inserting the UVC should have undertaken a formal training package for the insertion of central venous catheters. This should include an assessment of technical competence and awareness of potential complications
- When inserting a UVC each UVC should be withdrawn to a point at which it freely aspirates blood (to prevent malposition) and, after being secured in position, should be x-rayed to confirm that the position is acceptable.
- Following any new manual adjustment of UVC position, irrespective of how small the adjustment, a further radiograph must be obtained to verify the new position.
- The UVC position should also be noted on any subsequent x-ray done in the baby
- Umbilical catheters should be clearly labelled to distinguish arterial and venous catheters.
- Use of ultrasound to confirm ongoing position can be used where facilities and skill mix allows.<sup>27</sup>
- There is no evidence to support the practice of 'railroading' a second UVC alongside the one which is thought to be malpositioned. This practice is to be avoided since it may increase the risk of vessel trauma and consequent extravasation.
- There should be thorough contemporaneous documentation of each UVC insertion including indication, description of the catheter itself, number of attempts, length inserted, confirmation of blood aspiration, position on X-ray, and any adjustments subsequently made. The accepted position should be verified in

writing within 24 hours of insertion by a consultant neonatologist/consultant paediatrician or from a consultant radiologist's report.

- The need for continued retention/use of a UVC should be reviewed daily.
- On-going care of UVC should include regular review of catheter fixation and position, strict asepsis and minimising catheter access.
- Any clinical deterioration of a baby in whom a central venous catheter is present should raise the question of catheter-related complications, particularly infection, extravasation and tamponade
- Parents should always be informed about the use of central catheters at the earliest opportunity, although formal prior consent is impractical.

## 8) Description and Documentation of the Procedure

### Equipment

- Sterile pack for insertion of umbilical lines
- Chlorhexidine antiseptic solution for skin preparation
- Sterile water
- Umbilical catheter (single or double lumen, in size range 3.0 to 5.0 Fr)
- 10 mL syringe
- Yellow needle (21G)
- 0.9% sodium chloride – 10 mL ampoule
- Scalpel
- Sterile gloves and gown
- Suture
- Tape to secure the line in place with suture or umbilical catheter holder
- Umbilical tape

### Preparation

- Clean trolley surface.
- Wash hands.
- Assess the depth that the catheter needs to be inserted from one of the following two methods:

a)  $\{(3 \times \text{baby's weight in kg}) + 9\} / 2$  and add on cord stump length, (cm)<sup>17</sup>

b) Directly measure length from cord base to xiphisternum and add on cord stump length

- Position the infant and surrounding equipment so that the cord is accessible.
- Where possible, depending on the urgency of the procedure ensure that infant's temperature is at least 37°C before starting the procedure. Check that there is adequate output from the radiant heat source or incubator to keep the infant warm during the procedure.
- If the infant is particularly active and doesn't calm when the drapes are in place ensure that an assistant is on hand to contain and support the infant for the procedure. Consider use of sucrose if applicable.

### Physiological instability during insertion


- Closely observe the infant during and following the procedure for any deterioration
- Monitoring (ECG and oxygen saturation) should remain in place throughout the procedure
- If the infant is intubated, check the endotracheal tube is secure before commencing the procedure.
- If the infant is very preterm and is nursed in a polyethylene bag this should remain in place with a small incision made to the polyethylene bag over the umbilicus to provide access.

### Procedure

- Wash hands.
- Open the packaging of equipment with a non-touch technique.
- Wash hands thoroughly and dry.
- Put on gown and sterile gloves *in addition, hat & mask should be worn where this is local unit policy*
- Follow aseptic procedure principles.
- Draw up 10 mL of 0.9% sodium chloride into syringe and attach a three-way tap to the catheter. Flush through both the three-way tap (if used) and the catheter with the saline ensuring that there is no air in the system.
- Turn the three-way tap off or clamp the line to prevent any entry of air into the catheter to reduce the risk of air embolism whilst the catheter is being inserted.
- Clean cord and peri-umbilical area with chlorhexidine solution. Avoid excess application and any spillage around down to the buttocks as this may cause burns to

very preterm skin. Allow to dry, then remove with sterile water to prevent adverse skin reactions <sup>18,19</sup>.

- Holding cord clamp with sterile gauze, apply sterile drapes.
- Tie umbilical tape around the base of the cord tightly enough to minimise blood loss but loosely enough to allow the catheter to be passed through.
- Grasp the cord with the artery forceps and gently pull the cord upwards whilst you cut the underside of the forceps with a scalpel blade leaving 1-2cm of cord above the skin junction.
- Identify the vessels in the cord. The vein is a thin walled vessel, larger than the two arteries.
- Hold the stump with the toothed forceps and remove any visible clots.
- Place the tip of the catheter, held with forceps (see **Figure 1**) into the lumen of the vessel and gently advance to 5 cm into the vein. Remember that the vein goes up towards the heart unlike the arteries which descend first before looping upward; therefore, the catheter should be passed upwards.
- Turn the three-way tap so that the catheter is open to the syringe and apply gentle suction on the syringe.
- If there is smooth blood flow, continue to insert to the predetermined length and aspirate with syringe to verify blood return.



**Figure 1** Inserting catheter into umbilical vein (from 'Atlas of Procedures in Neonatology' <sup>2</sup>.)

- If no blood is aspirated at this point either advance the catheter 1 cm at a time or withdraw catheter 1 cm at a time until blood can be aspirated.
- If it is not easy to get blood back, rarely the catheter may be in a small blood vessel and have a clot in the tip – withdraw whilst maintaining suction, remove the clot and

reinsert the catheter. Note that UVCs placed during newborn resuscitation also require demonstration of adequate back sampling of blood before use.

- If there is any resistance and you cannot advance the catheter to the desired depth or there is a bobbing motion of the catheter, it may have entered the portal vein or be wedged in the intra-hepatic branch of the umbilical vein. The catheter will either must be retracted or removed and replaced. Some authors have described a technique involving insertion of another smaller bore catheter alongside which may then pass into the ductus venosus before removal of the original catheter<sup>21</sup>. However, there is insufficient evidence to support the technique of double cannulation at the present time. This practice may increase the risk of umbilical vein trauma and should therefore be avoided pending better supportive evidence for its safety<sup>1</sup>.
- Take blood samples as required and flush the line with saline.
- Secure the catheter using a technique that avoids tape being applied to the skin if possible e.g. suture and flag or Sulle securing device secured with a suture to the umbilicus<sup>22,23</sup>. Or use a colloid based umbilical catheter holder that will protect the skin e.g. Neobridge<sup>24</sup>. Use cavilon spray prior to using an adhesive on the skin. Once secured in place the catheter should not be advanced any further into the vein during any subsequent adjustments.
- Running 0.9% saline solution at 1.0 mL/hr through the catheter while awaiting X-ray confirmation of position may reduce the risk of the catheter blocking.
- Ensure that the correct position of the catheter is confirmed on AXR/CXR. The catheter should lie outside the cardiac silhouette ideally at T8-T9<sup>1</sup>. A UVC tip sited at or below T10 should only be use short term (if considered essential) as situation at or below this level carries a significantly higher risk of extravasation.<sup>1</sup> **Figures 2 and 3** illustrate the umbilical vein and its main anatomical relations.





Figure 2 Schematic diagram of umbilical and associated veins





**Figure 3 Anatomy of the umbilical and associated veins, with reference to external landmarks (from 'Atlas of Procedures in Neonatology' <sup>2</sup>)**

- There may be some bleeding from the umbilical vein because it is not a contractile vessel. Applying pressure may be enough to stop the oozing or the ligature around the bottom of the stump can be tightened slightly for a short while but released once oozing has stopped.
- Connect the catheter to the prescribed infusion.
- Clear away all equipment and ensure that any needles or scalpel are safely disposed of into a sharps bin.
- Record the procedure in the infants' medical notes using a dedicated UVC documentation label, such as the example in Figure 4., The UVC label should provide for the routine specific documented confirmation that there was blood sampling from the UVC at the point of catheter fixation.
- Regular nursing observations of the catheter insertion site and of the external anatomical site corresponding to the catheter's internal route should be undertaken (for example using a Visual Infusion Phlebitis "VIP" scoring system). It is recommended that as a minimum hourly inspection and documentation of the state of the umbilical catheter insertion site is recorded.

- Need for the UVC should be reviewed daily. The catheter should be removed as soon as it is no longer required, to prevent complications such as thrombosis and infection<sup>6,25,26</sup>.

**Insertion of UVC**

Date:..... Time:.....

Indication .....

Insertion length: use one of following methods

i)  $(\{3 \times \text{weight in kg}\} + 9) / 2 + \text{cord stump length. cm}$  or

ii) Directly measure length from cord base to xiphisternum and add on cord stump length

Length by formula: .....cm. Stump length: ..... cm

Actual insertion length: .....cm

Confirm that line sampled blood at insertion length:  (tick)

No. of attempts.....

Person undertaking catheter insertion: Name.....

Designation..... Signature .....

Date and time:..... Successful / Unsuccessful

---

Position on X-Ray: ..... Acceptable Yes or no

If no, state action taken:.....

Position on repeat X-Ray:..... Acceptable Yes or no

---

Sign below to verify line position confirmed as satisfactory by Consultant Neonatologist/paediatrician/radiologist (within 24 hours of insertion)

Name & Signature: Date & Time

**Fig 4: Example of umbilical venous catheterisation sticker for document casenotes.**

## References

1. British Association of Perinatal Medicine (2015) Use of Central Venous Catheters in Neonates A Framework for Practice. Revised 2018. London: BAPM. Available at: <http://www.bapm.org/publications/documents/guidelines/BAPM%20CVC%20final%20Dec15.pdf> (Last accessed: 21<sup>st</sup> August 2020)
2. MacDonald MG. (2002) Umbilical vein catheterisation. Ch 28. In: MacDonald MG, Ramasethu J. *Atlas of procedures in neonatology*. 3rd Ed. Lippincott Williams & Wilkins. Philadelphia.
3. Anderson, J. Leonard, D. Braner D.A.V, Lai, S. and Tegtmeyer, K. (2008) Videos in clinical medicine. Umbilical vascular catheterization. *The New England Journal of Medicine*. June; 359:(15);e18
4. Landers S, Moise AA, Fraley JK, Smith EO, Baker CJ. (1991) Factors associated with umbilical catheter-related sepsis in neonates. *American Journal of Disease in Childhood*. June;145(6):675-80.
5. Lam HS, Li AM, Chu WC, Yeung CK, Fok TF, Ng PC. (2005) Mal-positioned umbilical venous catheter causing liver abscess in a preterm neonate. *Biology of the Neonate*. March;88(1):54-6.
6. Kim JH, Lee YS, Kim SH, Lee SK, Lim MK, Kim HS. (2001) Does umbilical vein catheterization lead to portal venous thrombosis? Prospective US evaluation in 100 neonates. *Radiology*. 219:645-650.
7. Sehgal A, Cook V, Dunn M. (2007) Pericardial effusion associate with an appropriately placed umbilical venous catheter. *Journal of Perinatology*. May;27(5):317-9.
8. Department of Health. Review of the deaths of four babies due to cardiac tamponade associated with the presence of a central venous catheter. Department of Health, London, 2001. <http://tinyurl.com/nzc7wlx> (accessed 10th April 2016).
9. Pabalan MJ, Wynn RJ, Reynolds AM, Ryan RM, Youssfi M, Manja V, Lakshminrusimha S. (2007) Pleural effusion with parenteral nutrition solution: an unisial complication of an “appropriately” placed umbilical venous catheter. *American Journal of Perinatology*. November;24(10):581-5.
10. Dhya, A.K. Singh, B. Kumar, D. Awasthy, N. (2013) Broken umbilical vein Catheter as an embolus in a neonate- An unusual preventable complication. *Journal of Neonatal Surgery*. October; 2(4):45.
11. Onal EE, Saygili A, Koc E, Turkyilmaz C, Okumus N, Atalay Y. (2004) Cardiac tamponade in a newborn because of umbilical venous catheterisation: is correct position safe? November;14(11):953-6.
12. van Niekerk M, Kalis NN, van de Merwe PL. (1998) Cardiac tamponade following umbilical vein catheterisation in a neonate. *South African Medical Journal*. April;88(Suppl 2):C87-90.
13. Traen M, Schepens E, Laroche S, van Overmeire B. (2005) Cardiac tamponade and pericardial effusion due to umbilical venous catheterisation. *Acta Paediatrica*. May;94(5):626-8.
14. Sinha A, Fernanades CJ, Kim JJ, Fenrich AL, Encisco J. (2005) Atrial flutter following placement of an umbilical venous catheter. July;22(5):275-7.
15. Coley BD, Seguin J, Cordero L, Hogan MJ, Rosenberg E, Reber K. (1998) Neonatal total parenteral nutrition ascites from liver erosion by umbilical vein catheters. *Pediatric Radiology*. December;28(12):923-7.

16. Nakstad B, Naess PA, de Lange C, Schistad O. (2002) Complications of umbilical vein catheterisation: neonatal total parenteral nutrition ascites after surgical repair of congenital diaphragmatic hernia. *Journal of Pediatric Surgery*. August;37(8):E21.
17. Ahluwalia JS, Brain JL, Kelsall AW. (2005) Ch 45 Procedures and iatrogenic disorders. In: Rennie JM. *Robertson's Textbook of Neonatology*. Elsevier Churchill Livingstone. Philadelphia .
18. AWHONN and NANN (2001) *Evidence-based clinical practice guideline: Neonatal skin care*. Washington.
19. Reynolds PR, Banerjee S, Meek JH. (2005) Alcohol burns in extremely low birthweight infants: still occurring. *Archives of Disease in Childhood Fetal and Neonatal Edition*. January;90(1):F10.
20. GomellaTL (1994) Venous Access Ch 25. In: GomellaTL, Cunningham MD, Eyal FG. *Neonatology* 3rd Ed. Prentice-Hall. London. [IV]
21. Davies MW, Cartwright DW. (1998) Insertion of umbilical venous catheters past the ductus venosus using the double catheter technique. *Archives of Disease in Childhood Fetal and Neonatal Edition*. May;78(3):F234. [IV]
22. South M, Magnay A. (1988) Simple method for securing umbilical catheters. *Archives of Disease in Childhood*. July;63(7 Spec No):750-1. [IV]
23. Hindley DT, Lewis MA, Robinson MJ. (1994) Method for securing umbilical lines. *Archives of Disease in Childhood Fetal and Neonatal Edition*. January;70(1):F79-80. [IV]
24. Conway DP. (1998) Improved umbilical catheter anchoring device. *Neonatal Network*. August;17(5):65.
25. Rudzińska I, Gawecka A, Juszczak A, Kornacka MK. (2006) Central venous catheter colonization and catheter-related infection in preterm infants. *Journal of Pediatrics and Neonatology*. 3(2):NT39-42.
26. Butler-O'Hara M, Buzzard CJ, Reubens L, McDermott MP, DiGrazio W, Angio CT. (2006) A randomized trial comparing long-term and short-term use of umbilical venous catheters in premature infants with birth weights of less than 1251 grams. *Pediatrics*. July;118(1):e25-35.
27. Meinen, R.D., Bauer, A.S., Devous, K. et al. Point-of-care ultrasound use in umbilical line placement: a review. *J Perinatol* 40, 560–566 (2020).  
<https://doi.org/10.1038/s41372-019-0558-8>

**All Rights Reserved. The East of England Neonatal ODN withholds all rights to the maximum extent allowable under law. Any unauthorised broadcasting, public performance, copying or re-recording will constitute infringement of copyright. Any reproduction must be authorised and consulted with by the holding organisation (East of England Neonatal ODN).**

**The organisation is open to share the document for supporting or reference purposes but appropriate authorisation and discussion must take place to ensure any clinical risk is mitigated. The document must not incur alteration that may pose patients at potential risk. The East of England Neonatal ODN accepts no legal responsibility against any unlawful reproduction. The document only applies to the East of England region with due process followed in agreeing the content.**

## Exceptional Circumstances Form

Form to be completed in the **exceptional** circumstances that the Trust is not able to follow ODN approved guidelines.

Details of person completing the form:	
Title:	Organisation:
First name:	Email contact address:
Surname:	Telephone contact number:
Title of document to be excepted from:	
Rationale why Trust is unable to adhere to the document:	
Signature of speciality Clinical Lead:	Signature of Trust Nursing / Medical Director:
Date:	Date:
Hard Copy Received by ODN (date and sign):	Date acknowledgement receipt sent out:

Please email form to: [mandybaker6@nhs.net](mailto:mandybaker6@nhs.net) requesting receipt.

Send hard signed copy to: Mandy Baker  
 EOE ODN Executive Administrator  
 Box 93  
 Cambridge University Hospital  
 Hills Road Cambridge CB2 0QQ