

Joint Trust Guidelines for the Limping Child with no History of Trauma

A Clinical Guideline

| | |
|---|--|
| For Use in: | Children's Assessment Unit |
| By: | Medical and Nursing staff |
| For: | Children (0-16) presenting with a limp or acute lower limb pain but with no history of trauma |
| Division responsible for document: | Women / Children |
| Key words: | Limping, child, irritable, hip, septic, arthritis, osteomyelitis, limp |
| Name and job title of document author's: | Dr Julie Riechmann, Paediatric Junior Clinical Fellow; Dr Charlotte Lindsay, Emergency Medicine Registrar; Dr Aravind Shastri, Paediatric Consultant; Mr Anish Sanghrajka, Paediatric Orthopaedic Consultant |
| Name and job title of document author's Line Manager: | Dr Mary Anne Morris, Chief of Service for Paediatrics |
| Supported by: | Dr Sara Abdelgalil, Paediatric Consultant; Dr Vijayalakshmi Wardley, Paediatric Consultant; Miss Helen Chase, Paediatric Orthopaedic Consultant; Dr Abigail Reeve, Paediatric Consultant (JPUH). |
| Assessed and approved by the: | Clinical Guidelines Assessment Panel (CGAP) If approved by committee or Governance Lead Chair's Action; tick here <input checked="" type="checkbox"/> |
| Date of approval: | 02/06/2020 |
| Ratified by or reported as approved to (if applicable): | Clinical Safety and Effectiveness Sub-Board |
| To be reviewed before: This document remains current after this date but will be under review | 02/06/2023 |
| To be reviewed by: | Dr Aravind Shastri |
| Reference and / or Trust Docs ID No: | JCG0034 – Id 1235 |
| Version No: | 4.1 |
| Compliance links: (is there any NICE related to guidance) | No |
| If Yes - does the strategy/policy deviate from the recommendations of NICE? If so why? | N/A |

This guideline has been approved by the Trust's Clinical Guidelines Assessment Panel as an aid to the diagnosis and management of relevant patients and clinical circumstances. Not every patient or situation fits neatly into a standard guideline scenario and the guideline must be interpreted and applied in practice in the light of prevailing clinical circumstances, the diagnostic and treatment options available and the professional judgement, knowledge and expertise of relevant clinicians. It is advised that the rationale for any departure from relevant guidance should be documented in the patient's case notes.

The Trust's guidelines are made publicly available as part of the collective endeavour to continuously improve the quality of healthcare through sharing medical experience and knowledge. The Trust accepts no responsibility for any misunderstanding or misapplication of this document.

Quick Reference

Clinical Guideline for: The Limping Child with no history of trauma

Author/s: Dr Riechmann, Dr Lindsay, Dr Shastri, Dr Sanghrajka

Author/s title: See front sheet

Approved by: CGAP

Date approved: 02/06/2020

Review date: 02/06/2023

Available via Trust Docs


Version: 4.1

Trust Docs ID: 1235

Page 1 of 8

Joint Trust Guidelines for the Limping Child with no History of Trauma

Assessment of a child with an atraumatic limp



| Version Number | Date of Update | Change Description | Author |
|----------------|----------------|--|---|
| 4.1 | 02/06/2020 | Document is now a 3 trust document. Local supporter added: Dr Abigail Reeve, Paediatric Consultant (JPUH). | Dr Julie Riechmann Dr Charlotte Lindsay Dr Aravind Shastri Mr Anish Sanghrajka |
| | | | |

This is a Controlled Document

Printed copies of this document may not be up to date. Please check the hospital intranet for the latest version and destroy all previous versions.

Objective of Guideline

To promote thorough assessment and rational management of children with acute lower limb pain, limp or non-weight bearing, without a history of injury or trauma.

Rationale for the recommendations

A pre-guideline survey found that both paediatricians and orthopaedic surgeons were managing children presenting with a limp and a possible diagnosis of irritable hip or septic arthritis. The management varied in investigations undertaken, admission and follow-up arrangements. This guideline was developed following a review of the literature and agreement between specialties to rationalise the management of such children.

Major differential diagnoses of a child with a limp.

| Condition: | Typical features: |
|---|---|
| Septic arthritis (hip or other joint) | Any age. Most common <2yrs. Often painful (pseudo-paralysis), non-weight bearing, fever and unwell – these are not necessary. Reduced ROM of affected joint. CRP > 20 is likely to be associated with septic arthritis. |
| Osteomyelitis | Any age, similar features to septic arthritis BUT often more indolent presentation. Partial treatment with antibiotics common. Look for bone tenderness. In under 2's often co-exists with septic arthritis |
| Transient synovitis of the hip (Irritable hip) | 3-9 years. Post viral. Pain and limp, decreased ROM of hip but not as painful as septic arthritis |
| Fracture – non-accidental injury or unrecognised trauma | Take history carefully. Be alert to late presentations, inconsistencies. Toddler fracture – often minor fall resulting in undisplaced tibial fracture. Be aware of fractured fibula |
| Inflammatory arthritis (reactive, JIA, Lyme disease, HSP) | Joint swelling and heat (not detectable in hip). Decreased ROM but not as painful as septic arthritis. Longer history. Limping and pain and stiffness worse in morning/ after period of rest |
| Late presentation of Developmental Dysplasia | Always walked with limp. May have asymmetrical skin creases, shortened leg, look for a limitation of abduction in flexion |
| Perthes disease | 3-10 years, boys>girls. Limp with groin, thigh or knee pain. Decreased ROM with internal rotation of hip often reduced first |

Joint Trust Guidelines for the Limping Child with no History of Trauma

| | |
|--|--|
| Slipped upper femoral epiphysis (SUFE) | 8-15 years, boys>girls. Longer history limp, sudden minor trauma often worsens pain and leads to presentation, knee pain common, decreased ROM hip |
| Neoplasia - leukaemia | Night pain. General malaise. Weight loss, hepato-spleno megally, pallor, bruising |
| Neoplasia - Osteosarcoma | Pain, localised swelling (knee especially), often no history of systemic symptoms or night pain |
| Discitis | <5y/o Patient refusing to walk. Back pain. Loss of lumbar lordosis on examination |
| Neurological | Always consider neurological causes in a longstanding limp. |

Assessment

Initial assessment and documentation:

Pain - assess on pain scales according to age. Give appropriate analgesia
Normal observations – inform a doctor urgently if concerned about sepsis.
Weight (height if able to stand).

Medical assessment:

Key points of history

- Pain – site, severity, radiation, duration, exacerbating and relieving factors. Limp – similar detail.
- A history of preceding viral symptoms is often found in irritable hip.
- Preceding streptococcal sore throat / diarrhoeal illness in reactive arthritis.
- Fever height, duration, frequency.
- Recent antibiotic use may mask or partially treat a septic arthritis / osteomyelitis.
- Is the child considered to be generally well or unwell? Is s/he eating and drinking normally.
- Duration of symptoms – between 1 and 5 days associated with increased risk of infection.

Examination key points

- Is the child generally well or unwell?
- What is the gait and are they able to weight bear?
- Observe, palpate and move all bones and joints (look for heat, erythema, swelling, pain, restriction). Ensure you examine the spine.
- Severity of pain?
- Fever $\geq 38.5^{\circ}\text{C}$ is likely to be associated with increased risk of infection.
- Conduct a detailed neurology examination, including eliciting deep tendon reflexes (DTR). Remember, a child who is not weight bearing and has abnormal neurological findings like absent reflexes may have an underlying neurological cause to their limp.

Joint Trust Guidelines for the Limping Child with no History of Trauma

Follow algorithm for management.

Contact orthopaedic registrar on call on bleep 0996 if septic arthritis or osteomyelitis are suspected in an unwell patient.

For Outpatient referrals:

- *PRACC Clinic*: Discuss with and email CAU Consultant On Call or Night registrar
- *Rheumatology Clinic*: Make a referral using link on intranet (Electronic Forms)
- *Orthopaedic Clinic* in 7-10days: Email secretaries: Kelly.tilbury@nnuh.nhs.uk / Deanna.elliott@nnuh.nhs.uk. Also email paed ortho Consultant On call (switch are aware of names) that day: Helen.Chase@nnuh.nhs.uk / Anish.Sanghrajka@nnuh.nhs.uk / Graeme.Carlile@nnuh.nhs.uk

Medication doses as per BNF November 2019:

| Patient Age | Paracetamol Dose |
|-------------------------|---|
| 28 weeks - 32 weeks CGA | 20 mg/kg 1 st dose, then 10–15 mg/kg 8–12 hrly PRN, Max 30 mg/kg/day |
| 32 weeks CGA + above | 20 mg/kg 1 st dose, then 10–15 mg/kg 6–8 hrly PRN. Max 60 mg/kg/day |
| 1–2 months | 30–60 mg 8hrly, Max 60 mg/kg/day |
| 3–5 months | 60 mg 4–6 hrly, Max QDS |
| 6-23 months | 120 mg mg 4–6 hrly, Max QDS |
| 2-3 years | 180 mg mg 4–6 hrly, Max QDS |
| 4-5 years | 240 mg mg 4–6 hrly, Max QDS |
| 6–7 years | 240–250 mg 4–6 hrly, Max QDS |
| 8–9 years | 360–375 mg mg 4–6 hrly, Max QDS |
| 10–11 years | 480–500 mg mg 4–6 hrly, Max QDS |
| 12–15 years | 480–750 mg mg 4–6 hrly, Max QDS |
| 16–17 years | 0.5–1 g mg 4–6 hrly, Max QDS |

| Patient Age | Ibuprofen dose |
|-------------|--|
| 1-2 months | 5 mg/kg TDS-QDS |
| 3-5 months | 50 mg TDS; Max 30 mg/kg/day. |
| 6-11 months | 50 mg TDS-QDS, Max 30 mg/kg/day |
| 1-3 years | 100 mg TDS, Max 30 mg/kg/day |
| 4-6 years | 150 mg TDS, Max 30 mg/kg/day |
| 7-9 years | 200 mg TDS, Max 30 mg/kg/day. Max 2.4g/day |
| 10-11years | 300 mg TDS, Max 30 mg/kg/day. Max 2.4g/day |
| 12-17years | 300–400 mg TDS, Max 600 mg QDS |

Joint Trust Guidelines for the Limping Child with no History of Trauma

Upper Reference Range Limits for bloods:

- WCC >12
- ESR >20
- CRP >20

General Advice

Careful and full examination is imperative if clues concerning non-accidental injury, osteomyelitis and septic arthritis at any site or inflammatory arthritis are to be detected. Fully expose, palpate and move all bones / joints including the spine. Ensure full general examination is done including ENT and urine dip if febrile. Consider haemarthroses in child with excessive bruising. Developmental dysplasia of hip may present late with limp. If red flags are identified then organise bloods and necessary imaging concurrently.

X-Ray affected bones / joints (or whole of lower limb if very difficult to localise problem). If a fracture is present consider mechanism and any child protection issues. Refer to orthopaedics for fracture management or in suspected primary bone malignancies (bleep orthopaedic SpR on call on 0996).

ASO titre, anti-DNAse B and viral serology if reactive arthritis likely. ANA, autoantibody screen, immunoglobulins and rheumatoid factor are indicated if arthritis is likely but are not required urgently, or at first presentation.

The key is not to miss children with a septic hip joint since severe destruction of the joint can occur within 24 hours if not treated. If they are non-weight bearing despite analgesia and you suspect infection, inform the paediatric orthopaedic team early. If any doubt, investigate with blood tests and consider USS and aspiration / observation / admission / paediatric orthopaedic opinion. If any bony abnormality such as Perthes, SUFE or primary bone tumours are seen on X-ray refer to paediatric orthopaedic team.

Admission guidance

Any child who is not able to weight bear after appropriate analgesia should be admitted.

If a child is generally unwell (fever and/or significant pain and tenderness) they need investigation and likely admission for a period of observation. A paediatric orthopaedic opinion should be sought for any admitted child.

Parents should be given open access to phone the children's assessment unit and return to hospital in the next 2 weeks with the same problem. The parent information sheet at the end of this guideline should be given to them, with instruction to return if:

- The child is not better after 7 days of rest.
- The child develops a high temperature or is generally not well in himself or herself.
- The child is in more pain or is not able to put weight on their leg to walk.
- If they have any concerns they should phone 01603 289774.

Clinical Audit Standards derived from guideline

Joint Trust Guidelines for the Limping Child with no History of Trauma

- Initial point of contact should be paediatric on call team.
- Appropriate investigations when required.
- Appropriate referral to orthopaedics.
- Irritable hips given analgesia and advise sheet.

Monitoring of: Length of time in CAU, any delay in diagnosis and treatment of septic arthritis / osteomyelitis

Summary of development and consultation process undertaken before registration and dissemination

The authors listed above have developed the guideline and circulated for comment from paediatricians, orthopaedic surgeons, paediatric/MSK radiologists, nursing staff on CAU, A&E consultants, junior medical staff. Changes were made following these discussions.

Distribution list / dissemination method

CAU, A&E, Intranet
Trustdocs

Abbreviations

ANA – anti-nuclear antibody
AP – Antero-posterior
BC - Blood culture
CAU – Children’s Assessment Unit
CRP – C-reactive protein
DDH – Developmental Dysplasia of the Hip (previously known as congenital dysplasia of hip)
ENT – Ear, nose and throat
ESR – Erythrocyte sedimentation rate
FBC – Full blood count
GA- general anaesthetic
HSP- Henoch Schönlein purpura
JIA – Juvenile idiopathic arthritis
MSK - musculoskeletal
ROM – range of movement
SHO – senior house officer
SpR- specialist registrar
SUFE – slipped upper femoral epiphysis
USS – ultra-sound scan
WCC – white cell count
XR – X-Ray

References / source documents

Kinnaird TP, Beach RC. Fractured fibula can mimic irritable hip. *Arch Dis Child* 2003;**88**:167

Tuten HR, Gabos PG, Kumar SJ, Harter GD. The limping child: a manifestation of acute leukaemia. *J Pediatr Orthop* 1998;**18**:625-9

Joint Trust Guidelines for the Limping Child with no History of Trauma

Kocher MS, Zurakowski D, Kasser JR. Differentiating between septic arthritis and transient synovitis of the hip in children: an evidence-based clinical prediction algorithm. *J Bone & Joint Surg-Am* 1999;**81**:1662-70

Luhmann SJ, Jones A, Schootman M, Gordon JE, Schoenecker PL, Luhmann JD. Differentiation between septic arthritis and transient synovitis of the hip in children with clinical prediction algorithms. *J Bone Joint Surg Am* 2004;**86**:956-62

Fischer SU, Beattie TF. The limping child: epidemiology, assessment and outcome. *J Bone Joint Surg Br* 1999;**81**:1029-34

Kermond S, Fink M, Graham K, Carlin JB, Barnett P. A randomised clinical trial: should the child with transient synovitis of the hip be treated with non-steroidal anti-inflammatory drugs? *Ann Emerg Med* 2002;**40**:294-9

Beach RC. Minimally invasive approach to management of irritable hip in children. *The Lancet* 2000;**355**:1202-03

Jung ST, Rowe SM et al. Significance of laboratory and radiologic findings for differentiating between septic arthritis and transient synovitis of the hip. *J Pediatr Orthop* 2003;**23**:368-72

JaiKumar,Manoj Ramachandran, David Little,Michalis Zenios Pelvic Osteomyelitis in children, Journal of Paediatric Orthopaedics B 2010, **19**: 38-41

R.A.Delaney, B Leneham, L O’Sullivan, A.J.McGuinness, J.T.Street, Ir.Med.Sc : (2007)**176**:181-187

Markus Pa“a”kko“nen , Markku J. T. Kallio , Pentti E. Kallio , Heikki Peltola, Clin Orthop Relat Res (2010) 468:861–866

M Paakkonen, H Peltola, Management of a child with suspected acute septic arthritis, arch Dis Child 2012: 97: 287-292

National Institute for Health and Care Excellence. Clinical Knowledge Summaries: Acute Childhood Limp [Internet]. [London]: NICE; 2015 [updated 2015 Aug; cited 2020 Jan 13]. Available from: <https://cks.nice.org.uk/acute-childhood-limp#!topicSummary>