



**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

**Version and Document Control:**

<b>Version No.</b>	<b>Date of Update</b>	<b>Change Description</b>	<b>Author</b>
1	04/07/2014	Change of header and reference to joint hospital version	Akesh Dhrampal
2	18/10/2016	No clinical changes	Akesh Dhrampal
3	July 2020	Guidance on Direct Oral Anticoagulants (DOACs) added and references updated. Edoxaban added to the list of anticoagulants on the Epidural / Paravertebral / Interscalene brachial plexus Catheter Risk Assessment Tool in the Appendix on page 11.	Akesh Dhrampal

**This is a Controlled Document**

Printed copies of this document may not be up to date. Please check the hospital intranet for the latest version and destroy all previous versions.

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

**Quick Reference Guide for Regional Anaesthesia with:**

**a. Low Molecular Weight Heparin (LMWH): Dalteparin  $\leq$ 5000U, Tinzaparin  $\leq$  4500U or Enoxaparin  $\leq$  40mg S/C daily.**

If LMWH administered before block:	Wait at least 12 hours before block insertion.
If block insertion before LMWH:	Wait at least 4 hours before LMWH administration.
Epidural catheter removal:	Wait at least 12 hours since last dose and 4 hours before next dose of LMWH.

Administer LMWH at 18:00 so that at least 12 hours normally elapses before surgery the following day.
Removal of epidural catheters should occur between 10:00 and 14:00 to allow suitable intervals between doses of LMWH administered at 18:00. Epidural removal risk assessment tool (Appendix 2) should be used.

**b. Unfractionated heparin (UFH) 5000 IU 8 or 12 hourly**

If UFH administered before block:	Wait 4 hours before block insertion.
If block inserted before UFH:	Wait 1 hour before UFH administration.
Epidural catheter removal:	Wait 4 hours since last dose and 1 hour before next dose of UFH.

**Delay Venous thromboembolism (VTE) prophylaxis if:**

1. A hemorrhagic aspirate (i.e. a “bloody tap”) is encountered during the initial neuroaxial needle or epidural catheter placement, delay VTE prophylaxis by 24 hours.
2. When a neuroaxial technique is attempted but abandoned for general anesthesia, patients probably sustain excessive trauma to the epidural space. In this situation, it is better to wait at least 24 hours before initiating LMWH therapy.
3. Patients undergoing surgery in the late afternoon, where there is <4 hours between neuroaxial block insertion and routine LMWH administration time of 18:00, should have their VTE prophylaxis deferred until at least 4 hours have elapsed post epidural/spinal insertion.

**Therapeutic doses of both UFH and LMWH, markedly increase the risk of bleeding. Under these circumstances, neuroaxial anaesthesia should not be attempted. Occasionally patients may be therapeutically anticoagulated with UFH or LMWH after neuroaxial catheter insertion. Under no circumstances must these epidural catheters be removed while the patient is therapeutically anticoagulated.**

## Joint Trust Guideline for the Management of: Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs

Patients  $\leq 50$ kg see advice sheet ([Trustdocs ID: 1697](#)) for dose of Thromboprophylactic LMWH. Patients  $\leq 50$ kg who receive a conventional dose LMWH should have their epidural removed after 24 hours.

### Warfarin therapy:

For elective surgery warfarin should be stopped pre-operatively. ([Trustdocs ID: 1215](#))

INR must be  $\leq 1.5$ :

1. Within 24 hours of insertion of spinal/epidural needles and catheters.
2. Whenever the indwelling epidural catheter is in place.

Warfarin should **NOT** be given whilst an epidural catheter is in situ.

### c. Drugs with Antiplatelet activity: ([Trustdocs ID: 9836](#))

**Check indication for antiplatelet agent before stopping. If coronary artery stents present discuss with cardiologist.**

1. Aspirin, NSAIDs and dipyridamole:  
If used alone, appear to represent no added significant risk for the development of spinal hematoma in patients having epidural or spinal anesthesia.
2. Thienopyridine derivatives e.g. clopidogrel/prasugrel: Stop 7 days before surgery. Do not give whilst epidural in situ.
3. GP IIb/IIIa inhibitors (e.g. abciximab/Tirofiban): Contraindicated with neuroaxial analgesia techniques.

### Objective/s

To prevent the occurrence of vertebral canal haematomas when neuroaxial anaesthesia is administered to patients receiving drugs with anticoagulant and antiplatelet activity.

### Rationale

The incidence of VTE is increased markedly in association with surgery, trauma and immobilised medical patients. Therefore effective VTE prophylaxis is now a standard of care required in these patients<sup>7</sup>.

Neuroaxial anaesthesia (spinal or epidural) confer many advantages, including a significant reduction in the incidence of VTE and is commonly practised locally. Anticoagulants may increase the risk of bleeding from vertebral vessels damaged during the insertion of neuroaxial anaesthesia. This can lead to permanent neurological injury if a vertebral canal haematoma develops. Although such an event is devastating, it is an extremely rare occurrence and far more patients suffer from the effects of VTE.

Patients should have the combined benefits of VTE prophylaxis and neuroaxial anaesthesia, with minimal risk of vertebral canal haematoma formation. To this effect, awareness of anticoagulant status and careful coordination of the

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

administration of VTE prophylaxis and neuroaxial anaesthesia insertion is essential. The decision to institute neuroaxial anaesthesia must be made after assessing the risks versus benefits for each individual patient.

Prescribers need to be mindful of the fact that anticoagulants differ greatly in their ability to alter clotting, with different pharmacodynamic and pharmacokinetic properties. Co-morbidities (e.g. renal failure), and concurrent administration of drugs with anticoagulant/antiplatelet activity, may enhance the effects of VTE prophylaxis medication. Careful monitoring for spinal cord compression and prompt effective action in the event of neurological symptoms will enhance safety.

**Broad recommendations**

This guideline should be used in conjunction with Trust guideline CA2029/B17 for the management of adult patients receiving Epidural analgesia ([TrustDocs ID: 1911](#)) and guideline CA2060 for the management of adult patients on therapeutic anticoagulation who require elective surgery or an invasive procedure ([TrustDocs ID: 1215](#)).

- a) Low Molecular Weight Heparin (Dalteparin  $\leq$  5000U, Tinzaparin  $\leq$  4500U or Enoxaparin  $\leq$  40mg S/C daily).

If LMWH administered before block:	Wait at least 12 hours before block insertion.
If block insertion before LMWH:	Wait at least 4 hours before LMWH administration.
Epidural catheter removal:	Wait at least 12 hours since last dose and 4 hours before next dose of LMWH.

For the majority of elective patients these criteria will be met by the use of the following dosing schedule.

Administer LMWH at 18:00 so that at least 12 hours normally elapses before surgery the following day.
Removal of epidural catheters should occur between 10:00 and 14:00 to allow suitable intervals between doses of LMWH administered at 18:00. Epidural removal risk assessment tool (Appendix 2) should be used.

- b. Unfractionated Heparin (5000 U S/C 8 or 12 hourly).

If UFH administered before block:	Wait 4 hours before block insertion.
If block inserted before UFH:	Wait 1 hour before UFH administration.
Epidural catheter removal:	Wait 4 hours since last dose and 1 hour before next dose of UFH.

New to these guidelines is the recommendation that VTE prophylaxis should be delayed by 24 hours if:

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

1. A hemorrhagic aspirate (i.e. a “bloody tap”) is encountered during the initial neuroaxial needle or epidural catheter placement.
  
2. When a neuroaxial technique is attempted but abandoned for general anesthesia, patients probably sustain excessive trauma to the epidural space. In this situation, it is better to wait at least 24 hours before initiating LMWH therapy.
  
3. Patients undergoing surgery in the late afternoon, where there is <4 hours between neuroaxial block insertion and routine LMWH administration time of 18:00, should have their VTE prophylaxis deferred until at least 4 hours have elapsed post epidural/spinal insertion.

**Therapeutic doses of both UFH and LMWH, markedly increase the risk of bleeding. Under these circumstances, neuroaxial anaesthesia should not be attempted. Occasionally patients may be therapeutically anticoagulated with UFH or LMWH after neuroaxial catheter insertion. Under no circumstances must these epidural catheters be removed while the patient is therapeutically anticoagulated.**

c. Regional Anaesthesia and Warfarin therapy:

Patients receiving warfarin therapy should have this discontinued  $\geq 5$  days before the planned surgical procedure (see [Trustdocs ID: 1215](#) for the management of adult patients on therapeutic anticoagulation who require elective surgery or an invasive procedure).

An INR should be checked within 24 hours of the procedure and a value of  $\leq 1.5$  is considered safe for spinal/epidural insertion. Warfarin therapy is **contraindicated** whilst an epidural catheter is in situ.

d. Regional Anaesthesia and drugs with Antiplatelet activity:

Aspirin, NSAIDs and dipyridamole: If used alone these agents appear to represent no added significant risk for the development of spinal hematoma in patients having epidural or spinal anesthesia. At present, there are no specific concerns as to the timing of single-shot or catheter techniques in relationship to the dosing of NSAIDs, postoperative monitoring, or the timing of neuraxial catheter removal.

Thienopyridine derivatives (e.g. clopidogrel/prasugrel): Consensus management is based on labeling precautions and surgical, interventional cardiology/radiology experience. Clopidogrel should be discontinued 7 days before neuroaxial blockade is attempted.

**Antiplatelet agents must not be stopped in patients who have coronary artery stents insitu without discussion with a cardiologist.**

These agents should not be recommenced in a patient whilst an indwelling epidural catheter is insitu.

## Joint Trust Guideline for the Management of: Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs

GP IIb/IIIa inhibitors (e.g. abciximab), exert potent antiplatelet effects. Accordingly neuroaxial techniques should be avoided until platelet function has recovered. GP IIb/IIIa antagonists are contraindicated within four weeks of surgery. Should one need to be administered in the postoperative period (following a neuroaxial technique), the indwelling epidural catheter should be removed first.

In summary, the concurrent use of other medications which may affect clotting mechanisms may increase the risk of bleeding and hence spinal haematomas if a neuroaxial block is sited. Extra vigilance is required and the patient monitored closely for signs of cord compression (Appendix 1).

d. Direct Oral Anticoagulants (DOACs). Also known as Non-Vitamin K oral anticoagulants (NOAC's) or Target-Specific Oral Anticoagulants (TSOAC's) (Trustdocs ID: 8752)

Non-vitamin K oral anticoagulants (dabigatran (Pradaxa<sup>®</sup>), rivaroxaban (Xarelto<sup>®</sup>), apixaban (Eliquis<sup>®</sup>) and edoxaban (Lixiana<sup>®</sup>)) are options for the prevention of stroke and systemic embolism in people with non-valvular atrial fibrillation (AF). Generally can be stopped 2 days before operation (longer if deranged renal function). Should not be given while epidural catheter insitu.

f. Removal of epidural catheters

To ensure safe removal of epidural catheter the trust epidural risk assessment tool should be completed before removal. This is available on all surgical wards where epidural catheters are in routine use. See appendix 2.

g. Diagnosis and management of vertebral haematoma

If a haematoma is suspected the guidance in Appendix 1 should be followed.

### Clinical audit standards

Adherence to recommended time intervals between:

- a. Needle insertion and VTE prophylaxis administration.
- b. Epidural catheter removal and VTE prophylaxis administration.

Completion of epidural removal risk assessment tool.

### Summary of development and consultation process undertaken before registration and dissemination

The authors listed above drafted this guideline on behalf of the anaesthetic department who has agreed the final content. During its development it has been circulated for comment to representatives of the general surgery, orthopaedic/spinal surgery, obstetric & gynaecological surgery, haematology and surgical ward nursing staff.

In July/August 2020 this document was reviewed by the Thrombosis and Thromboprophylaxis Committee and the Clinical Guidelines Assessment Panel.

# Joint Trust Guideline for the Management of: Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs

## Distribution list/dissemination method

Anaesthetists, Surgeons, Ward nursing staff, Clinical guidelines folder, Intranet.

## References/source documents

1. Wildsmith JAW, McClure JH. Anticoagulation drugs and central nerve blockade. *Anaesthesia* 1991;46:613-4
2. American society of Regional Anaesthesia. Consensus Statement 2002. [www.asra.com](http://www.asra.com)
3. American College of Chest Physicians 7<sup>th</sup> Guidelines on Thromboprophylaxis. *Chest* 2004
4. Tryba M. European Practice Guidelines: Thromboembolism prophylaxis and regional anaesthesia. *Regional Anaesthesia and Pain Medicine* 1998;23:178-182
5. Wu CL. Regional Anaesthesia and anticoagulation. *J Clinical Anaesthesiology* 2001;13:49-58
6. Scottish Intercollegiate Guideline Network. Prophylaxis of Venous thromboembolism 2002. [www.sign.ac.uk](http://www.sign.ac.uk)
7. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism. [NICE guideline \[NG89\]](#) Published date: 21 March 2018 Last updated: 13 August 2019.



### **Vertebral/Spinal Haematomas complicating Neuroaxial anaesthesia**

Although spinal haematomas in this setting is rare, it nevertheless could have a potentially devastating neurological outcome if not recognized early.

The following factors have been implicated in increasing the risk of spinal haematoma formation:

1. Frail, elderly females.
2. Associated haemostatic disorder.
3. Technically difficult insertion requiring multiple attempts at multiple levels.
4. Bloody tap i.e. blood in needle or epidural catheter.
5. Inappropriate doses or timing of LMWH.
6. Combination of drugs with anticoagulant/antiplatelet effects.
7. Removal of epidural catheter: 30-60% of clinically important haematomas occur after catheter removal. Most occur within 24 hours of catheter removal, but may occur up to 6 days later.

If neuroaxial anaesthesia was difficult or associated with haemorrhage, this should be documented and the patient monitored particularly closely for evidence of cord compression. Moreover, it is the recommendation of these guidelines that VTE prophylaxis should then be delayed for 24 hours.

#### **Detection:**

The symptoms and signs of spinal cord compression closely resemble the effects of a dense spinal or epidural block, which compounds the diagnostic accuracy. Therefore to assess the neurological function in the postoperative period, it is advisable to use infusion of opioids or low concentrations of local anaesthetic.

Medical staff, nursing staff and the patient must be made aware of what symptoms and signs to lookout for (patient information leaflet) so that the alarm can be raised. New onset of one or more of the following is suggestive of spinal cord compression.

1. Increasing motor weakness despite a constant epidural infusion rate.
2. Increasing sensory weakness despite a constant epidural infusion rate.
3. Cauda Equina signs i.e. Bladder and bowel incontinence.
4. Back pain which is often of sudden onset and “excruciating”.
5. Radicular pain due to irritation of nerve roots by blood.

Golden rules to be aware of:

- The patient **MUST** be able to move something on his legs, even if just dorsiflexing his toes or foot, the **WHOLE** time an epidural is running in the postoperative period.
- Movement should return 2-3 hours after a local anaesthetic bolus.

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

- Thoracic epidurals should be associated with little or no lower limb motor weakness.

**Treatment:**

This is a medical emergency, as spinal cord compression is likely to lead to permanent neurological injury if not surgically decompressed within 8 hours. The sooner the problem is recognised the sooner the surgery can be undertaken. If spinal haematoma is suspected stop epidural infusion and contact the on-call anaesthetist who will then liaise with the Consultant Spinal Surgeon and Radiologist to arrange an MRI followed by surgery if indicated.'

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

**B**

**Appendix 2**

<b>Pain Assessment and Observation frequency guidance:</b>		Addressograph
Epidural PCA:	Hourly for 8 hours 2 hourly for 48 hours 4 hourly thereafter	
Oral/IM/SC analgesia	Once per shift 1 hour after analgesia	

	<b>Pain</b>	<b>Sedation</b>	<b>Nausea</b>	<b>Catheter site</b>	<b>Leg Movement</b>
0	No pain	Alert	None	Clean	Full movement
1	Mild pain on movement	Mild: Easy to rouse	Mild nausea	Skin reddening, No induration	Weakness of plantar flexion
2	Moderate pain on movement	Easy to rouse, often drowsy	Nausea and retching	Skin induration No discharge	Weakness of hip flexion
3	Severe pain on movement	Somnolent. Difficult to rouse	Vomiting	Discharge from catheter	Unable to move legs

	Date/ time or 24 clock	00	01	02	03	04	05	06	07	08	09	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
Pain	3																									
	2																									
	1																									
	0																									
Sedation	3																									
	2																									
	1																									
	0																									
Nausea	3																									
	2																									
	1																									
	0																									
PCA	Tries																									
	Good																									
	Resp																									
	Total																									
Epidural	Rate																									
	Site																									
	Leg Mvmt																									
	Resp																									
	Total																									
Signature																										

**Joint Trust Guideline for the Management of:  
Regional Anaesthesia (Spinal and Epidurals) in patients receiving Venous  
Thromboembolism Prophylaxis, Anticoagulant and Antiplatelet Drugs**

**B**

**Epidural / Paravertebral / Interscalene brachial plexus Catheter Risk  
Assessment Tool**

**Appendix 3**

**Patient Name:**

**Hospital Number:**

**Before removing an epidural / paravertebral / interscalene brachial plexus catheter  
please answer all the questions below:**

Question	Yes	No
<b>A. Heparin</b>		
1. Is the patient prescribed Dalteparin (Fragmin) greater than 5,000 units or Tinzaparin (Innohep) dose greater than 4,500 units or Enoxaparin (Clexane) greater than 40 mg		
2. Has <u>any</u> heparin (Dalteparin, Tinzaparin, Enoxaparin or Unfractionated Heparin -sc or iv) been given in the last 12 hours?		
3. If the patient weighs less than 50 kgs have they received <b>any</b> heparin in the last <b>24</b> hours?		
<b>B. Other anti-coagulant or anti-platelet Drugs</b>		
4. Has Warfarin been given within the past 2 days?		
5. For patients who received Warfarin in the last week – is the INR greater than 1.5 <b>on the day of removal</b> ?		
6. Has the patient received Rivaroxaban within the last 18 hours?		
7. Has the patient received Fondaparinux or Dabigatran within the last 36 hours?		
8. Has Clopidogrel or Prasugrel been given within the past 7 days?		
9. Is the patient taking Aspirin doses greater than 300mg per day?		
10. Is the patient taking any other anticoagulants or anti-platelet drugs**?		
<b>C. Early Warning Score</b>		
11. At time of assessment does the patient trigger the Early Warning Score (EWS)?		
<b>If the answer is YES to any of the questions, DO NOT remove the epidural, PVB or brachial plexus catheter Seek advice from the Pain Team on Bleep 0571 or the On-call Anaesthetist</b>		
If unable to remove (any YES response) please document the name of the person contacted and advice given:		
<b>Signature:</b>	<b>Date</b>	<b>Time:</b>
<b>Epidural / paravertebral / interscalene brachial plexus catheter removed?</b>		
<b>Print Name:</b>	<b>Signature:</b>	
<b>Date:</b>	<b>Time:</b>	
<b>Patient Information Leaflet given?</b>	<b>Yes <input type="checkbox"/></b>	<b>Signature:</b>
<b>Dose Post Removal Advice: Drugs may be restarted after the following time intervals</b>		
4 hours: Unfractionated Heparin, Warfarin 4-6 hours: Dalteparin, Tinzaparin, Enoxaparin Next day: Rivaroxaban, Dabigatran, Fondaparinux, Clopidogrel, Aspirin, Prasugrel		
** Other anti-coagulant and anti-platelet agents include (this is not an exhaustive list): Abciximab, Acenocoumarol, Apixaban, Bivalirudin, Danaparoid, Edoxaban, Epoprostenol, Idraparinux, Phenindione, Ticagrelor, Tirofiban		
<b>For further information see Adult Patients Receiving Epidural Analgesia <a href="#">Trustdocs ID: 1191</a></b>		