Speech and Language Therapy Department

Having a PEG tube



 Speech and Language Therapy Department Having a PEG tube

 Author/s: Speech and Language Therapy Department

 Approved by: PIF
 Date approved: 03/01/2024

 Available via Trust Docs
 Version: 2.3
 Trust Docs ID: 11646

On the day of the Procedure

- Do not eat and drink by mouth for 6 hours before.
- You can have water up to 2 hours before if you can swallow safely.
- Your doctor will tell you which medications you can take.
- If you are being fed by tube this will be stopped 6 hours before.
- You can have water through your tube up to 2 hours before.

What happens when you have a PEG?

- It takes about **30 minutes**. •
- You will have the feeding tube in hospital.
- You will have a local anaesthetic and sedative.
- You will be awake but sleepy.
- A doctor passes the **tube** through your **mouth** and **food pipe** down to the **stomach**.
- A small cut is made in the skin in your stomach.
- Then the tube is **pulled out** from the **stomach**.





Will it hurt?

- You may have **pain** in your **stomach** after the tube is put in.
- You will be given painkillers to help.

Speech and Language Therapy Department Having a PEG tube Author/s: Speech and Language Therapy Department Approved by: PIF Date approved: 03/01/2024 Trust Docs ID: 11646



What happens next

- First you will have water through your • tube.
- Liquid feed will then be started. The dietitian will talk to you about your feed.
- The dietitian will teach you how to use the •

Will I be able to eat and drink?

The Speech and Language Therapist will test your swallow.



Speech and Language Therapy Department Having a PEG tube Author/s: Speech and Language Therapy Department Approved by: PIF Date approved: 03/01/2024

Review date: 03/01/2027 Page 3 of 4



