

## HAEMATOLOGY H.A009 - THROMBOPHILIA ADVICE

### Role of thrombophilia screening:

Venous thromboembolism (VTE) is a multifactorial disease. There is a thrombotic threshold for everyone, which if exceeded will cause thrombosis. The threshold is altered throughout life by environmental and lifestyle factors. Detectable thrombophilic defects are only **one** of the known risk factors for thrombosis and should not be acted upon in isolation. Every patient with VTE should be assessed for clinical risk factors first - e.g. obesity, recent surgery, immobility, increasing age, malignancy – as some may be modifiable. In most instances thrombophilia screening contributes little to future management.

### General points about thrombophilia screens:

- For the vast majority of patients presenting with VTE, thrombophilia testing is of limited benefit.
- The finding of heritable thrombophilia should not influence decisions regarding the intensity of anticoagulation.
- Duration of anticoagulation after VTE is mainly determined by clinical factors estimating the risk of recurrent VTE, and the risk of bleeding associated with anticoagulation.
- The advice regarding thromboprophylaxis to prevent future thrombosis remains unchanged.
- Thrombophilia results should not be used in isolation to determine if it is safe to prescribe oestrogen containing contraceptives, HRT or other medications known to carry a thrombotic risk (see specific Gynaecology department guidance in these situations).

### Who to consider for screening?

- Selected individuals from thrombosis prone families can be considered for testing. Care should be taken before testing to consider how the results will be of benefit. In almost all patients, the results do not influence current or subsequent management. People with a strong family history of thrombosis should be considered at increased risk, irrespective of the results of a thrombophilia screen.
- Pregnant women with a history of VTE, or who have a family history of VTE in a first degree relative. This may help make decisions about thromboprophylaxis in pregnancy.
- As part of investigations for pregnancy loss.
- Individuals who require investigation for suspected antiphospholipid syndrome (request lupus anticoagulant/anticardiolipin antibodies only).
- Selected patients after unprovoked VTE who are planning to stop anticoagulation if the results would change management. Testing for antiphospholipid syndrome (APS) and testing for heritable thrombophilia in patients who also have a first degree relative with a history of VTE could be considered.

---

This is a controlled document and must be read in conjunction with all NNUH NHS Trust Policies and Procedures.

It is the responsibility of the user to ensure that they are aware of the current issue and printed copies (including blank forms) can only be deemed current at the time of printing (04/11/2022 15:20). Please notify any changes required to the document approver.

Title: Thrombophilia

Page 1 of 4

Review due date: 3<sup>rd</sup> February 2023

Document reference: H.A009.v7 Trust Document ref : 4648

- Selected patients with unprovoked VTE and risk factors for APS. Testing for APS can be considered if the result would change choice of anticoagulant. Features in VTE patients which can be suggestive of APS are: history of SLE or autoimmune disease, livedo reticularis, prolonged APTT prior to anticoagulation, recurrent VTE, VTE at unusual sites, history of arterial thrombosis without risk factors, thrombocytopenia, recurrent miscarriage/still birth/severe pre-eclampsia).

### When is screening not advised?

- Patients with VTE where anticoagulation is being continued indefinitely should not be tested for hereditary thrombophilia. The result of a hereditary thrombophilia test will not change management (see above for suspected APS).
- Patients with provoked VTE.
- Asymptomatic patients with no personal or family history of VTE.
- Children under 16. Genetic screening of children under 16 is not advised without good clinical indication and genetic counselling.
- Results of previous inherited thrombophilia screen available. The genetic tests should not be repeated; clotting based assays are as accurate as methodology allows and should only be repeated if advised to do so by the laboratory or haematology.
- Arterial thrombosis. Patients with arterial thrombosis should not be tested for heritable thrombophilia. Where the cause of arterial thrombosis is not known, consideration can be given to testing for antiphospholipid syndrome.

### Types and timing of thrombophilia screens:

There are 2 types of screen:

#### Heritable thrombophilia screen

- Protein C (clotting based)
- Protein S (clotting based)
- Antithrombin (clotting based)
- Factor V Leiden (genetic test)
- Prothrombin gene mutation (genetic test)

#### Acquired (not hereditary) thrombophilia Screen

- Lupus anticoagulant (clotting based)
- Anticardiolipin antibodies (ELISA based)

For a full thrombophilia screen send 3 x blue, 1 yellow and 1 purple vacutainer if all tests required. Otherwise select specific tube for test (clotting based = blue; genetic = purple; ELISA = yellow)

**Clotting based** assays are affected by anticoagulants (heparin, warfarin, DOACs - apixaban, edoxaban, dabigatran and rivaroxaban) and by acute thrombosis, and therefore should not be performed at the time of an acute clot, or whilst a patient is anticoagulated. The clotting based tests should be performed 4-6 weeks after anticoagulation stops. Genetic tests are **not** affected by

---

This is a controlled document and must be read in conjunction with all NNUH NHS Trust Policies and Procedures.

It is the responsibility of the user to ensure that they are aware of the current issue and printed copies (including blank forms) can only be deemed current at the time of printing (04/11/2022 15:20). Please notify any changes required to the document approver.

Title: Thrombophilia

Page 2 of 4

Review due date: 3<sup>rd</sup> February 2023

Document reference: H.A009.v7 Trust Document ref : 4648

the above and can be performed at any time. If testing on anticoagulants (excluding warfarin) is deemed beneficial, discuss with the laboratory regarding options for anticoagulant removal from the sample to permit testing.

### Interpretation of results

Acquired causes of reduced Protein C, Protein S and antithrombin levels occur more frequently than hereditary deficiency. Hereditary deficiency should only be diagnosed after exclusion of acquired causes and confirmed on repeat testing by a haematologist.

Acquired causes of low Protein C, Protein S and antithrombin levels are listed below:

**Protein C:** Vitamin K antagonists (warfarin), liver disease, DIC

**Protein S:** Pregnancy, combined oral contraceptive, vitamin K antagonists (warfarin), vitamin K deficiency, liver disease, sickle cell disease, acute chickenpox infection

**Antithrombin:** Acute thrombosis, heparin therapy, liver disease, nephrotic syndrome, asparaginase therapy

### Lupus anticoagulant and anticardiolipin antibodies

Positive results can be found in normal healthy individuals. A diagnosis of antiphospholipid syndrome should only be made in patients fulfilling the specific clinical diagnostic criteria (thrombotic or obstetric) and persistently positive tests (2 positive results at least 12 weeks apart).

### Estimate of risks of venous thromboembolism:

- lifetime overall risk of VTE 5%
- Incidence of VTE (increases with age) 1:1000 person years
- Incidence of VTE in women age < 40 yrs not on combined OCP 1 :10,000 person years
- Incidence of VTE in women age <40yrs on combined OCP 4 :10,000 person years
- Incidence of VTE in pregnancy 1-2:1000 pregnancies
- Incidence of VTE in women aged 50-59 yrs not on HRT 7-9:10,000 person years
- Incidence of VTE in women aged 50-59 yrs taking HRT 18-23:10,000 person years

### Estimates of prevalence of thrombophilic abnormalities and relative risks of thrombosis:

| thrombophilic abnormality               | prevalence in population | relative risk for first VTE | relative risk for recurrent VTE |
|---|--------------------------|-----------------------------|---------------------------------|
| antithrombin deficiency                 | 0.02%                    | 5-10                        | 1.9-2.6                         |
| protein C deficiency                    | 0.2%                     | 4-6.5                       | 1.4-1.8                         |
| protein S deficiency                    | 0.03% - 0.13%            | 1-10                        | 1.0-1.4                         |
| factor V Leiden mutation (heterozygous) | 3% - 7%                  | 3-5                         | 1.4                             |
| prothrombin gene mutation               | 0.7% – 4%                | 2-3                         | 1.4                             |

NB. The absolute risk of thrombosis for any given patient will depend on age and other thrombotic risk factors. Absolute risks can be more informative when counselling patients regarding their individual risk of thrombosis.

## References:

Does thrombophilia testing help in the clinical management of patients? S Middeldorp et al  
British Journal of Haematology Volume 143: 321–335, November 2008  
<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2141.2008.07339.x/full>

Clinical guideline for testing for heritable thrombophilia. T Baglin et al. British Journal of  
Haematology Volume 149: 209–220, April 2010  
<http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2141.2009.08022.x/full>

Addendum to British Society for Haematology guidelines on Investigation and Management of  
Antiphospholipid syndrome 2012. Published Jan 2020.  
<https://onlinelibrary.wiley.com/doi/abs/10.1111/bjh.16308>

NICE guidance NG158: Venous thromboembolic disease: diagnosis, management and  
thrombophilia testing. March 2020.  
<https://www.nice.org.uk/guidance/ng158/chapter/Recommendations#thrombophilia-testing>