

## East of England Perinatal Network

### Clinical Guideline: Enteral Feeding – Vitamin supplementation

**Authors:** Lynne Radbone, Principal Paediatric Dietitian

**For use in:** EoE Neonatal Units  
Guidance specific to the care of neonatal patients.

**Used by:** Medical staff, Neonatal Nurse Practitioners, Dietitians, Nursery Nurses


**Key Words:** vitamins, enteral feeding, Abidec, Folic acid

**Date of Ratification:**

**Review due:**

**Registration No: NEO-ODN-2018-18**

**Approved by:**

Neonatal Clinical Oversight Group	
Clinical Lead Mark Dyke	

**Ratified by Eastern Perinatal Board:**

<b>Ratified 13<sup>th</sup> December 2018</b>	<b>Review December 2021</b>
---	---------------------------------

**Audit Standards:**

#### Audit points

All babies born at <34 weeks gestation:

Receiving unfortified expressed breast milk or donor breast milk are given 0.6ml Abidec and 50 micrograms folic acid daily (the latter until term).

Receiving fortified expressed breast milk, preterm formula, term formula, specialist formulas or NEPDF are given 0.6ml Abidec daily.

Who are discharged exclusively breast feeding receive 0.6ml Abidec daily until 1 year corrected age.

Who are discharged on term, specialist formulas or NEPDF receive 0.6ml Abidec daily until 6 months corrected age.

## 1. Introduction

The third trimester of pregnancy is a time of rapid nutrient accretion, peak bone formation and the time when vitamin stores are laid down, infants born prematurely therefore have lower stores and higher requirements for vitamins than those born at term. Additionally low levels of specific enzymes and carrier proteins lead to lower absorption and transportation of some nutrients.

A number of vitamins have been studied in relation to prematurity, notably Vitamin A, Vitamin D, Vitamin E and Folic Acid. These form the basis of supplementation recommendations.

The gestation below which additional vitamins are required is unclear and supplementation practice has in the past varies across the EOE Network. International guidelines from Koletzko (2014) provide recommendations for vitamin intakes in ELBW and VLBW infants and ESPGHAN (2010) for infants <1800g but neither make any delineation by degree of prematurity.

As infants born >33 weeks are most likely to establish breast feeding quicker than more premature infants, by consequence they will correct any nutrient deficit within a shorter period of time. For these reasons these guidelines should apply at the earliest to infants <34 weeks gestation.

Vitamin supplements should commence once an enteral feed volume of >100ml/kg/day has been achieved alongside a reciprocal reduction in aqueous and lipid PN. Vitamins are contained within the lipid fraction of PN so care should be taken to ensure parenteral lipid provision is 5ml/kg/day or less before commencing enteral vitamin supplements. In contrast, consideration should be given to earlier enteral supplementation if PN weaning practice involves the cessation of lipid prior to that of aqueous PN.

Vitamin supplementation is necessary for many infants born prematurely. The evidence base for the exact requirement is limited for most vitamins, and although the quantities required are extremely small they are all essential to many basis life processes. As such they should be included in any enteral supplementation guidance. (1)

The following guidelines are based on a combination of careful analysis of the vitamin content of available feeds and formulas, available evidence and best practice.

The available evidence used to support these recommendations can be found in Appendix 2.

## 2. Vitamin supplementation regimen

Guidelines apply to all preterm infants <34 weeks gestation on full enteral feeds according to the milk they are receiving. Care should be given to changes in milk types and modifications made to the vitamin supplement regimen as appropriate.

**Dalivit should not be used as an interchangeable alternative for Abidec due to its far higher vitamin A content. (Appendix 2)**

Milk type	Vitamin supplement	Dose
<b>In the Neonatal Unit</b>		
Unfortified breast milk	Folic Acid Abidec	50micrograms/day 0.6ml/day
EBM+ SMA Breast Milk Fortifier (BMF) EBM + Nutriprem Human Milk Fortifier (HMF)	Abidec	0.6ml/day
Nutriprem 1 Hydrolysed Nutriprem SMA Gold Prem 1	Abidec	0.6ml/day
Term formula Specialist formulas Nutrient dense term formulas	Abidec	0.6ml/day
<b>At time of discharge</b>		
SMA Gold Prem 2 Nutriprem 2	Abidec	0.6ml/day until 6 months corrected age
Breast feeding	Abidec	0.6ml/day until 1 year corrected age
Term formula Specialist formulas Nutrient dense term formulas	Abidec	0.6ml/day until 6 months corrected age

### **Vitamin supplementation in infants with conjugated hyperbilirubinaemia..**


Infants with an increasing conjugated bilirubin >50micromol/l may have a degree of fat malabsorption, thereby indicating a need for additional supplements of fat soluble vitamins. The following supplementation regimen is suggested for these infants within the neonatal unit.

**All other vitamin supplementation should be stopped when this regimen is implemented.**


Preparation	Dose	Provision
Dalivit	0.6ml/day	5000 units vitamin A 400 units vitamin D
Alpha- tocopherol acetate suspension	10mg/kg/day (2)	14.9 units vitamin E / kg
Phytomenadione	1mg daily (3)	




3. Algorithm for Vitamin supplementation: **Nutriprem Range of Feeds**




Maximum feed volumes should not exceed **180ml/kg** when using EBM + Nutriprem HMF, Nutriprem 1 or Hydrolysed Nutriprem.



Algorithm for Vitamin supplementation: **SMA Range of Feeds**



Maximum feed volumes should not exceed **150ml/kg** when using EBM + SMA Breast Milk Fortifier or SMA Gold Prem 1 preterm formula.



#### 4. After discharge

The duration of increased vitamin requirements in preterm infants is unclear, though until 5kg has been suggested in the US.(4) The preterm infant at term is not comparable to the infant born at term having started out with, and accumulated, significant nutritional deficits in the early stages of life (5).

UK current practice is to continue supplementation until 6-12 months actual age and/or established on a good mixed diet. The Department of Health recommend that all term infants from 6 months of age who are breastfed or receiving less than 500ml formula per day should receive supplementary vitamin A, D and C as a matter of course (6).

NEPDF are prescribable until a maximum of 6 months corrected age after which term formula or a "follow on" formula are recommended. At this point either 0.3ml Abidec or 5 drops of the Healthy Start Vitamin preparation should be commenced as per the Department of Health National Guidelines.

Healthy Start Vitamin Drops are not available on prescription. They are available free of charge to those families who qualify for Healthy Start Vouchers or can be purchased over the counter at selected pharmacies and retailers. Infants who are not eligible for the Healthy Start Scheme, or where local purchase options are not available should receive Abidec (not Dalivit) on prescription from their General Practitioner.

It is recommended that a standard letter or leaflet outlining vitamin supplement requirements are included in discharge planning packages for both parents and General Practitioners.

## Appendix 1 Composition of available vitamin supplements

**Dalivit should not be used as an interchangeable alternative for Abidec or Healthy Start Vitamins due to its high vitamin A content.**

	Vit A	Vit D	Vit C	Vit B1	Vit B2	Nic Acid	Vit B6
<b><u>Abidec</u></b>	IU micrograms	IU micrograms	mg	mg	mg	mg	mg
<b>0.3ml</b>	666 200	200 5	20	0.2	0.4	4.0	0.4
<b>0.6ml</b>	1333 400	400 10	40	0.4	0.8	8.0	0.8
<b><u>Dalivit</u></b>							
<b>0.3ml</b>	2500 750	200 5	25	0.5	0.2	2.5	0.25
<b>0.6ml</b>	5000 1500	400 10	50	1.0	0.4	5.0	0.5
<b><u>Healthy Start</u></b>							
<b>5 drops</b>	660 200	280 7	20				
<b>10 drops</b>	1320 400	560 14	40				

## Appendix 2 - Evidence to support recommendations:

### Vitamin A

Vitamin A is required for growth and differentiation of epithelial tissues, including the lungs and the retina. Preterm infants have low vitamin A levels at birth, which appears to persist at discharge (7) and may contribute to an increased risk of developing Chronic Lung Disease. A Cochrane review undertaken in 2011 concluded that large doses of vitamin A given intramuscularly showed a small decrease in oxygen dependency in infants at 36 weeks, but only to infants born < 1000g.(8) Supplementation of vitamin A may reduce the incidence of Retinopathy of Prematurity (ROP), Intra Ventricular Haemorrhage (IVH) and Necrotising Enterocolitis (NEC) whereas excess vitamin A can raise intracranial pressure, cause skin and mucosal membrane changes and vomiting.

The most recent recommendations for vitamin A requirements in preterm infants are 400 -1,100 microgram RE/kg/day or 1,330 – 3,330 IU/kg/day (9), this figure, published in 2014 is unchanged from the recommendations made by ESPGHAN in 2010 (10). Higher oral doses of 5000units/day do not seem to show any clinical benefit (11).

The requirements post discharge are unknown though low levels have been reported in infants until 6 months corrected age who were discharged on term formulas, whereas normal levels were identified in infants fed NEPDF to 2 months corrected age(12).



## Vitamin D

Vitamin D is essential for the absorption of calcium and phosphorus and is therefore vital in bone formation. Supplementation is of no benefit if there are inadequate supplies of these two minerals, though its exact mechanism is unknown. Bronner et al (13) showed that calcium absorption in low birth weight infants was directly proportional to the calcium intake and independent of daily vitamin D supplementation. In contrast Devlin et al (14,15) showed that calcium absorption increases from 50% to 71% when AGA preterm infants <1500g were fed banked human milk alone or supplemented with vitamin D without calcium fortification, demonstrating that vitamin D affected calcium absorption rates.

Preterm infants are able to hydroxylate vitamin D so do not need the active form (16).

Supplementation with excess active vitamin D may cause calcium resorption of the bone so should only be considered where there is clear biochemical deficiency or poor absorption eg. significant cholestatic liver disease.

Historically 400 IU/day of vitamin D has been considered adequate for optimal absorption in the presence of sufficient calcium, phosphorus and magnesium (17). Koo demonstrated that 800 IU/day is no better than 400 IU/day, but that 200 IU/day was inadequate for the prevention of osteoporosis and rickets of prematurity (18). In 2010 ESPGHAN recommended 800 -1000 IU/day (9) based on the prevalence of vitamin D deficiency in pregnant women (19) and the international consensus to increase circulating vitamin D levels in the general population (20), however the most recent recommendations suggest that 400 IU/day is adequate in the presence of sufficient quantities of calcium, phosphorus and magnesium, but that provision should increase up to 1000 IU/day if there is a likelihood of maternal depletion (21)

## B vitamins and Folic Acid

Recommended levels for most B vitamins are provided by routine vitamin supplementation or formula composition, however Folic acid deficiency (in the form of growth retardation, anaemia and small intestine morphology) can be expected in the preterm infant not fed fortified breast milk or preterm formula due to poor intrauterine stores, rapid growth and low levels of folic acid in preterm breast milk. Folic acid is not present in standard vitamin supplements so needs to be supplemented separately.

Unfortified breast milk contains approx 3micrograms/100ml so does not meet the current recommendations of 35-100micrograms /kg/day(9). Recommendations for Folic acid are based on low plasma and red cell folate levels in preterm formula fed infants without Folic acid supplementation as compared to those given a supplement of 50microgram/day who had levels comparable to breast fed term infants(22).

## Vitamin E

Vitamin E is a biological antioxidant with a role in the prevention of haemolytic anaemia and may protect against Bronchopulmonary dysplasia (BPD), ROP and IVH. A Cochrane Review stated that "vitamin E supplementation in preterm infants reduces the risk of intra cranial haemorrhage but increases the risk of sepsis."(23) a second study concluded that vitamin E reduces the risk of severe retinopathy and blindness in those studied but it increases the risk of sepsis. It has also been associated with a higher risk of NEC.(24)

Evidence for supplementation of vitamin E is conflicting, however as minimum requirements for vitamin E are met from both human and formula milk there is little clear evidence to support routine supplementation.(24)

## References

1	Greene H, Hambidge K, Schanler R, Tsang R. Guidelines for the use of vitamins, trace elements, calcium, magnesium, and phosphorus in infants and children receiving total parenteral nutrition: report of the Subcommittee on Pediatric Parenteral Nutrient Requirements from the Committee on Clinical Practice Issues of the American Society for Clinical Nutrition.[erratum appears in Am J Clin Nutr 1989 Jun;49(6):1332]. <i>American Journal of Clinical Nutrition</i> 1988;48(5):1324-42.
2	British National formulary for children (Nov 2018) <a href="https://bnfc.nice.org.uk/drug/alpha-tocopherol.html">https://bnfc.nice.org.uk/drug/alpha-tocopherol.html</a>
3	British National formulary for children (Nov 2018) <a href="https://bnfc.nice.org.uk/drug/phytomenadione.html">https://bnfc.nice.org.uk/drug/phytomenadione.html</a>
4	Groh-Wargo S, Thompson M, Hovasi Cox J. Nutritional care for high risk newborns 3 <sup>rd</sup> edition. <i>Precept Press 2000</i> .
5	Embleton NE, Pang N, Cooke RJ. Postnatal Malnutrition and Growth Retardation: an inevitable consequence of current recommendations for preterm infants? <i>2001 Pediatrics, 107, 270-273</i> .
6	Dept of Health. Weaning and The Weaning Diet, Report on Health and Social Subjects 45 <i>HMSO: Stationery Office, 1994</i> .
7	Mactier H, Galloway P, Hamilton R, Weaver L. Inadequacy of IV vitamin A supplementation of extremely preterm infants? <i>The Journal of Pediatrics</i> 2005;146(6):846-47.
8	Darlow B, Graham P. Vitamin A supplementation to prevent mortality and short and long-term morbidity in very low birthweight infants. <i>Cochrane Database of Systematic Reviews</i> 2011;10:CD000501.
9	Leaf, A. and Z. Lansdowne, <i>Vitamins - conventional uses and new insights</i> , in <i>Nutritional Care of Preterm Infants: Scientific Basis and Practical Guidelines</i> .2014, Karger. p. 153-166.
10	ESPGHAN. Enteral Nutrient Supply for Preterm Infants: Commentary from European Society for Paediatric Gastroenterology, Hepatology, and Nutrition Committee on Nutrition. <i>JPGN</i> 2010;50:1-9.
11	Wardle SP, Hughes A, Chen S, Shaw NJ. Randomised controlled trial of oral vitamin A supplementation in preterm infants to prevent chronic lung disease. <i>Archives of Disease in Childhood Fetal &amp; Neonatal Edition</i> 2001;84(1):F9-F13.
12	Peebles JM, Carlson SE, Werkman SH, Cooke RJ. Vitamin A status of preterm infants during infancy. <i>Am J Clin Nutr</i> 1991;53(6):1455-59.
13	Bronner F, Salle BL, Putet G et al. Net calcium absorption in premature infants: results of 103 metabolic balance studies. <i>Am J Clin Nutr</i> 1992;56:1037-44.
14	Devlin EE, Lopez V, Levy E et al. Developmental expression of calcitriol receptors, 9 kilodalton calcium binding protein and calcidiol 24 hydroxylase in human intestine. <i>Pediatr Res</i> 1996;40:664-70.
15	Delvin EE, Lopez V, Levy E et al. Calcitriol differentially modulates mRNA encoding receptors and calcium binding protein 0kDa in human fetal jejunum. <i>Biochem Biophys Res Commun</i> 1996;224:544-8.
16	Koo WWK, Tsang RC, Mineral requirements of low birth weight infants. <i>Journal of the American College of Nutrition</i> . 1991;10:474.
17	Tsang R, Uauy R, Koletzko B, Zlotkin S. <i>Nutrition of the Preterm Infant: Scientific Basis and Practical Guidelines 2nd Ed</i> : Digital Educational Publishing Inc, 2005.
18	Koo WW, Krug-Wispe S, Neylan M, Succop P, Oestreich AE, Tsang RC. Effect of three levels of vitamin D intake in preterm infants receiving high mineral-containing milk. <i>Journal of Pediatric Gastroenterology &amp; Nutrition</i> 1995;21(2):182-9.

19	Holmes VA, Barnes MS, Alexander HD, McFaul P, Wallace JMW. Vitamin D deficiency and insufficiency in pregnant women: a longitudinal study. <i>British Journal of Nutrition</i> 2009;102(06):876-81.
20	Vieth R, Bischoff-Ferrari H, Boucher BJ, Dawson-Hughes B, Garland CF, Heaney RP, et al. The urgent need to recommend an intake of vitamin D that is effective. <i>Am J Clin Nutr</i> 2007;85(3):649-50.
21	Mimouni, F.B., D. Mandel, and R. Lubetzky, <i>Calcium, phosphorus, magnesium and vitamin D requirements of the preterm infant</i> , in <i>Nutritional care of preterm infants: scientific basis and practical guidelines</i> . 2014, Karger. p. 146-147.
22	Ek J, Behncke L, Halvorsen KS, Magnus E. Plasma and red cell folate values and folate requirements in formula-fed premature infants. <i>European Journal of Pediatrics</i> 1984;142(2):78
23	Brion LP, Bell EF & Ragheveer TS. Vitamin E supplementation for prevention of morbidity and mortality in preterm infants. <i>Cochrane Database Syst Rev</i> , 4. 2003
24	Brion LP, Bell EF & Ragheveer TS. Variability in the dose of intravenous vitamin E given to very low birth weight infants. <i>J Perinatol</i> 2005; 25, 139-142.

**All Rights Reserved. The East of England Perinatal Network withholds all rights to the maximum extent allowable under law. Any unauthorised broadcasting, public performance, copying or re-recording will constitute infringement of copyright. Any reproduction must be authorised and consulted with by the holding organisation (East of England Perinatal Network).**

**The organisation is open to share the document for supporting or reference purposes but appropriate authorisation and discussion must take place to ensure any clinical risk is mitigated. The document must not incur alteration that may pose patients at potential risk. The East of England Perinatal Network accepts no legal responsibility against any unlawful reproduction. The document only applies to the East of England region with due process followed in agreeing the content.**

## Exceptional Circumstances Form

Form to be completed in the **exceptional** circumstances that the Trust is not able to follow ODN approved guidelines.

Details of person completing the form:	
Title:	Organisation:
First name:	Email contact address:
Surname:	Telephone contact number:
Title of document to be excepted from:	
Rationale why Trust is unable to adhere to the document:	
Signature of speciality Clinical Lead:	Signature of Trust Nursing / Medical Director:
Date:	Date:
Hard Copy Received by ODN (date and sign):	Date acknowledgement receipt sent out:

Please email form to: [mandybaker6@nhs.net](mailto:mandybaker6@nhs.net) requesting receipt.

Send hard signed copy to: Mandy Baker  
EOE ODN Executive Administrator  
Box 93  
Cambridge University Hospital  
Hills Road  
Cambridge CB2 0QQ

